

# Influence of climate and land use changes on recent trends of soil erosion rates within the Russian Plain

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

## Abstract

© 2018 John Wiley & Sons, Ltd. The Russian Plain (within the Russian Federation) occupies an area of approximately  $3.5 \times 10^6$  km<sup>2</sup>. Water erosion is the main land degradation factor within the Russian Plain. Previous quantitative assessments of erosion rates for the entire area of the Russian Plain were undertaken in the 1980s. The application of erosion models and analysis of different factor dynamics allow for the evaluation of the mean annual total soil losses and erosion rates for the post-Union of Soviet Socialist Republics period, as well as the determination of the trends of erosion rates and soil losses for different landscape zones, for 1980 and 2012. The significant reductions of cultivated land area in all of the landscape zones after 1991 are the main reason behind the 46% reduction in the total annual soil losses in 2012 compared with 1980. The most significant decrease in the soil erosion rate for cultivated lands (from 7.3 Mg ha<sup>-1</sup> yr<sup>-1</sup> in 1980 to 4.1 Mg ha<sup>-1</sup> yr<sup>-1</sup> in 2012) was identified in the forest zone of the Russian Plain. Soil erosion rates slightly decreased in the forest-steppe zone and increased in the steppe zone (from 3.9 to 4.6 Mg ha<sup>-1</sup> yr<sup>-1</sup>). We attribute the decreasing erosion rates in the forest and forest-steppe zones to the reduction of surface runoff during snowmelt. The increased use of perennial grasses in crop rotations promoted the reduction of erosion rates in the forest zone. The increasing erosion rates in the steppe zone were attributed to the increasing frequency of heavy rainstorms.

<http://dx.doi.org/10.1002/ldr.3061>

## Keywords

climate change, land use change, Russian Plain, soil erosion rate, USLE

## References

- [1] Audry, S., Schäfer, J., Blanc, G., & Jouanneau, J.-M. (2004). Fifty-year sedimentary record of heavy metal pollution (Cd, Zn, Cu, Pb) in the Lot River reservoirs (France). *Environmental Pollution*, 132, 413–426. <https://doi.org/10.1016/j.envpol.2004.05.025>
- [2] Auerswald, K., Fiener, P., & Dikau, D. (2009). Rates of sheet and rill erosion in Germany—A meta-analysis. *Geomorphology*, 111, 182–193. <https://doi.org/10.1016/j.geomorph.2009.04.018>
- [3] Brazier, R. (2004). Quantifying soil erosion by water in the UK: A review of monitoring and modelling approaches. *Progress in Physical Geography*, 28, 340–365. <https://doi.org/10.1191/0309133304pp415ra>
- [4] Cerdan, O., Govers, G., Le Bissonnais, Y., Van Oost, K., Poesen, J., Saby, N., ... Dostal, T. (2010). The rate and spatial distribution of soil erosion in Europe. *Geomorphology*, 122, 167–177. <https://doi.org/10.1016/j.geomorph.2010.06.011>

- [5] Cerdan, O., Le Bissonnais, Y., Govers, G., Lecomte, V., van Oost, K., Couturier, A., ... Dubreuil, N. (2004). Scale effect on runoff from experimental plots to catchments in agricultural areas in Normandy. *Journal of Hydrology*, 299, 4–14. <https://doi.org/10.1016/j.jhydrol.2004.02.017>
- [6] de Vente, J., & Poesen, J. (2005). Predicting soil erosion and sediment yield at the basin scale. *Scale issues and semi-quantitative models*. *Earth-Science Reviews*, 71, 95–125. <https://doi.org/10.1016/j.earscirev.2005.02.002>
- [7] Dotterweich, M., Rodzik, J., Zgłobicki, W., Schmitt, A., Schmidtchen, G., & Bork, H.-R. (2012). High resolution gully erosion and sedimentation processes, and land use changes since the Bronze Age and future trajectories in the Kazimierz Dolny area (Nałęczów Plateau, SE-Poland). *Catena*, 95, 50–62. <https://doi.org/10.1016/j.catena.2012.03.001>
- [8] Du, P., & Walling, D. E. (2012). Using Pb measurements to estimate sedimentation rates on river floodplains. *Journal of Environmental Radioactivity*, 103, 59–75. <https://doi.org/10.1016/j.jenvrad.2011.08.006>
- [9] Edwards, W. M., & Owens, I. B. (1991). Large storm effects on the total soil erosion. *Journal of Soil and Water Conservation*, 46, 75–78.
- [10] Estel, S., Kuemmerle, T., Alcántara, C., Levers, C., Prishchepov, A., & Hostert, P. (2015). Mapping farmland abandonment and recultivation across Europe using MODIS NDVI time series. *Remote Sensing of Environment*, 163, 312–325. <https://doi.org/10.1016/j.rse.2015.03.028>
- [11] Fiener, P., Neuhaus, P., & Botschek, J. (2013). Long-term trends in rainfall erosivity—Analysis of high resolution precipitation time series (1937–2007) from Western Germany. *Agricultural and Forest Meteorology*, 171–172, 115–123. <https://doi.org/10.1016/j.agrformet.2012.11.011>
- [12] Golosov, V. N. (2006). *Erosion and deposition processes in river basins of cultivated plains*. Moscow: GEOS (in Russian).
- [13] Golosov, V. N., Belyaev, V. R., Kuznetsova, Y. S., Markelov, M. V., & Shamshurina, E. N. (2008). Response of a small arable catchment sediment budget to introduction of soil conservation measures. In *Sediment dynamics in changing environments* (Vol. 325) (pp. 106–113). IAHS Publication.
- [14] Golosov, V. N., Belyaev, V. R., & Markelov, M. V. (2013). Application of Chernobyl-derived Cs fallout for sediment redistribution studies: Lessons from European Russia. *Hydrological Processes*, 27, 807–821. <https://doi.org/10.1002/hyp.9470>
- [15] Golosov, V. N., Belyaev, V. R., Markelov, M. V., & Kislenko, K. S. (2010). Overbank sedimentation rates on the flood plains of small rivers in Central European Russia. In *Sediment dynamics for a changing future* (Vol. 337) (pp. 129–136). IAHS Publ.
- [16] Golosov, V. N., Belyaev, V. R., Markelov, M. V., & Shamshurina, E. N. (2012). Specifics of sediment redistribution within a small arable catchment during different periods of its cultivation (Gracheva Loschina catchment, the Kursk region). *Geomorphology RAS*, 1, 25–35. (in Russian) <https://doi.org/10.15356/0435-428-2012-1-25-35>
- [17] Golosov, V. N., Belyaev, V. R., Shamshurina, E. N., & Kuznetsova, J. S. (2014). Use of an integrated approach for assessing soil redistribution in the river Vorobzha basin. In *Proceedings FAO-IAEA international symposium on managing the soil for food security and climate change adaptation and mitigation* (pp. 107–112). Rome: FAO.
- [18] Golosov, V. N., Gennadiev, A. N., Olson, K. R., Markelov, M. V., Zhidkin, A. P., Chendev, Y. G., & Kovach, R. G. (2011). Spatial and temporal features of soil erosion in the forest-steppe zone of the East-European plain. *Eurasian Soil Science*, 44, 794–801. <https://doi.org/10.1134/S1064229311070064>
- [19] Gonzalez-Hidalgo, J. C., Batalla, R. J., Cerdà, A., & De Luis, M. (2009). Contribution of the largest events to suspended sediment transport across the USA. *Land Degradation & Development*, 21, 83–91. <https://doi.org/10.1002/ldr.897>
- [20] Hanel, M., Pavlásková, A., & Kyselý, J. (2016). Trends in characteristics of sub-daily heavy precipitation and rainfall erosivity in the Czech Republic. *International Journal of Climatology*, 36, 1833–1845. <https://doi.org/10.1002/joc.4463>
- [21] Intergovernmental Panel on Climate Change (2007). *Climate change 2007: The physical science basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press. <http://www.ipcc.ch/ipccreports/ar4-wg1.htm>
- [22] International Union of Soil Sciences Working Group WRB (2006). *World reference base for soil Resources* (2nd ed.). Rome: World Soil Resources Reports No. 103. FAO.
- [23] Ivanov, M. A., Prishchepov, A. V., Golosov, V. N., Zalyaliev, R. R., Efimov, K. V., Kondrateva, A. A., ... Ionova, Y. K. (2017). Changes of cropland area in the river basins of the European part of Russia for the period 1985–2015 years, as a factor of soil erosion dynamics. *Current Problems in Remote Sensing of the Earth from Space*, 14(6), 149–157. (in Russian) <https://doi.org/10.21046/2070-7401-2017-14-6-149-157>
- [24] Kireeva, M. B., Frolova, N. L., Winde, F., Dzhamalov, R. G., Rets, E. P., Povalishnikova, E. S., & Pahomova, O. M. (2016). Low flow on the rivers of the European part of Russia and its hazards. *Geography, Environment, Sustainability*, 9(4), 33–47. <https://doi.org/10.24057/2071-9388-2016-9-4-33-47>

- [25] Kosmas, C., Danalatos, N., Cammeraat, L. H., Chabart, M., Diamantopoulos, J., Farand, R., ... Vacca, A. (1997). The effect of land use on runoff and soil erosion rates under Mediterranean conditions. *Catena*, 29, 45–59. [https://doi.org/10.1016/S0341-8162\(96\)00062-8](https://doi.org/10.1016/S0341-8162(96)00062-8)
- [26] Krasilnikov, P., Makarov, O., Alyabina, I., & Nachtergaele, F. (2016). Assessing soil degradation in northern Eurasia. *Geoderma Regional*, 7, 1–10. <https://doi.org/10.1016/j.geodrs.2015.11.002>
- [27] Krasnov, S. F., Dobrovolskaya, N. G., & Litvin, L. F. (2001). Spatial and temporal aspects of the rainfall erosivity evaluation. *Soil Erosion and Channel Processes*, 13, 8–17. (in Russian)
- [28] Larionov, G. A. (1993). Water and wind erosion: Main features and quantitative assessment. Moscow: Izdatelstvo Moskovskogo Universiteta. (in Russian)
- [29] Larionov, G. A., Dobrovolskaya, N. G., Krasnov, S. F., Liu, B. Y., & Nearing, M. A. (1998). Theoretical-empirical equation of topography factor for a statistical model of soil erosion by water. *Soil Erosion and Channel Processes*, 11, 25–44. (in Russian)
- [30] Litvin, L. F., Zorina, Y. F., Sidorchuk, A. Y., Chernov, A. V., & Golosov, V. N. (2003). Erosion and sedimentation on the Russian Plain; Part 1, Contemporary processes. *Hydrological Processes*, 17, 3335–3346. <https://doi.org/10.1002/hyp.1390>
- [31] Lyuri, D. I., Goryachkin, S. V., Karavaeva, N. A., Denisenko, E. A., & Nefedova, T. G. (2010). Dynamics of agricultural lands of Russia in XX century and post-agrogenic restoration of vegetation and soils. Moscow: GEOS (in Russian).
- [32] Maracchi, G., Sirotenko, O., & Bindi, M. (2005). Impacts of present and future climate variability on agriculture and forestry in the temperate regions: Europe. *Climate Change*, 70, 117–135. <https://doi.org/10.1007/s10584-005-9166-7>
- [33] Markelov, M. V., Golosov, V. N., & Belyaev, V. R. (2012). Changes in the sedimentation rates on the floodplains of small rivers in the central Russian Plain. *Vestnik of Moscow State University, Seriya 5, Geography*, 5, 70–76. (in Russian)
- [34] Milanova, E. V., Lioubimtseva, E. Y., Tcherkashin, P. A., & Yanvareva, L. F. (1999). Land use/cover change in Russia: Mapping and GIS. *Land Use Policy*, 16, 153–159. [https://doi.org/10.1016/S0264-8377\(99\)00011-3](https://doi.org/10.1016/S0264-8377(99)00011-3)
- [35] Mueller, E. N., & Pfister, A. (2011). Increasing occurrence of high-intensity rainstorm events relevant for the generation of soil erosion in a temperate lowland region in central Europe. *Journal of Hydrology*, 411, 266–278. <https://doi.org/10.1016/j.jhydrol.2011.10.005>
- [36] Nearing, M. A., Pruski, F. F., & O'Neal, M. R. (2004). Expected climate change impacts on soil erosion rates: A review. *Journal of Soil and Water Conservation*, 59, 43–50.
- [37] Øygarden, L. (1996). Erosion and surface runoff in small agricultural catchments. In *Erosion and sediment yield: Global and regional perspectives* (Vol. 236) (pp. 283–294). IAHS Publ.
- [38] Øygarden, L., Deelstra, J., Lagzdins, A., Bechmann, M., Greipsland, I., Kyllmar, K., ... Iital, A. (2014). Climate change and the potential effects on runoff and nitrogen losses in the Nordic-Baltic region. *Agriculture, Ecosystems & Environment*, 198, 114–126. <https://doi.org/10.1016/j.agee.2014.06.025>
- [39] Panagos, P., Borrelli, P., Poesen, J., Ballabio, C., Lugato, E., Meusburger, K., ... Alewell, C. (2015). The new assessment of soil loss by water erosion in Europe. *Environment Science & Policy*, 54, 438–447. <https://doi.org/10.1016/j.envsci.2015.08.012>
- [40] Park, H., Sherstiukov, A. B., Fedorov, A. N., Polyakov, I. V., & Walsh, J. E. (2014). An observation-based assessment of the influences of air temperature and snow depth on soil temperature in Russia. *Environmental Research Letters*, 9, 1–7. <https://doi.org/10.1088/1748-9326/9/6/064026>
- [41] Petelko, A. I., Golosov, V. N., & Belyaev, V. R. (2007). Experience of design of system of counter-erosion measures. In *Proceedings of the 10th international symposium on river sedimentation*, vol. 1, Moscow: 311–316.
- [42] Poesen, J., Nachtergaele, J., Verstraeten, G., & Valentin, C. (2003). Gully erosion and environmental change: Importance and research needs. *Catena*, 50, 91–133. [https://doi.org/10.1016/S0341-8162\(02\)00143-1](https://doi.org/10.1016/S0341-8162(02)00143-1)
- [43] Porto, P., Walling, D. E., La Spada, C., & Callegari, G. (2016). Validating the use of Cs measurements to derive the slope component of the sediment budget of a small rangeland catchment in Southern Italy. *Land Degradation and Development*, 27, 798–810. <https://doi.org/10.1002/ldr.2388>
- [44] R Core Team (2016). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.r-project.org/>
- [45] Razuvaev, V. N., Apasova, E. G., Martuganov, R. A., Steurer, P., & Vose, R. (1993). Daily temperature and precipitation data for 223 U.S.S.R. Stations. ORNL/CDIAC, Numerical data package – 040. Oak Ridge, Tennessee, USA: Oak Ridge National laboratory.
- [46] Renard, K. G., & Freimund, J. R. (1994). Using monthly precipitation data to estimate the R-factor in the revised USLE. *Journal of Hydrology*, 157, 287–306. [https://doi.org/10.1016/0022-1694\(94\)90110-4](https://doi.org/10.1016/0022-1694(94)90110-4)
- [47] Rosstat (2015). Russian Federal Service of State Statistics. Moscow, Russia. Retrieved from <http://www.gks.ru>

- [48] Rysin, I., Grigoriev, I., Zaytseva, M., Golosov, V., & Sharifullin, A. (2017). Long-term monitoring of gully erosion in Udmurt Republic, Russia. In Integrating monitoring and modelling for sediment dynamics (Vol. 375) (pp. 1–4). Proc. IAHS. <https://doi.org/10.5194/piahs-375-1-2017>
- [49] Scholz, G., Quinton, J. N., & Strauss, P. (2008). Soil erosion from sugar beet in Central Europe in response to climate change induced seasonal precipitation variations. *Catena*, 72, 91–105. <https://doi.org/10.1016/j.catena.2007.04.005>
- [50] Sidorchuk, A., Litvin, L., Golosov, V., & Cherhysh, A. (2006). European Russia and Byelorussia. In J. Boardman, & J. Poesen (Eds.), *Soil Erosion in Europe*. Chapter 1.8. Chichester, England: John Wiley & Sons. <https://doi.org/10.1002/0470859202.ch8>
- [51] Sidorchuk, A. Y., & Golosov, V. N. (2003). Erosion and sedimentation processes on the Russian Plain, II: The history of erosion and sedimentation during the period of intensive agriculture. *Hydrological Processes*, 17, 3347–3358. <https://doi.org/10.1002/hyp.1391>
- [52] Smetanová, A., Kožuch, M., & Čerňanský, J. (2009). The land use changes in 20th century and their geomorphological implications in lowland agricultural area (Voderady, Trnavská tabuľa Table Plain, Slovakia). *Geomorphologica Slovaca et Bohemica*, 2, 57–63.
- [53] *Soil and Water Conservation Advances in the United States*. (2010). Zobeck T. M., & Schillinger W. F. (Eds.), Soil Science Society of America, Special Publication 60, Madison, WI: USA.
- [54] *Soil Erosion in Europe*. (2006). Boardman J., & Poesen J. (Eds.), John Wiley & Sons: Chichester, England. <https://doi.org/10.1002/0470859202>
- [55] Trimble, S. W., & Crosson, P. (2000). Land use—US soil erosion rates—Myth and reality. *Science*, 289, 248–250. <https://doi.org/10.1126/science.289.5477.248>
- [56] Van Rompaey, A., Krasa, J., & Dostal, T. (2007). Modelling the impact of land cover changes in the Czech Republic on sediment delivery. *Land Use Policy*, 24, 576–583. <https://doi.org/10.1016/j.landusepol.2005.10.003>
- [57] Walling, D. E. (2009). The impact of global change on erosion and sediment transport by rivers: Current progress and future challenges. Paper for the UNESCO-IHP International Sediment Initiative (ISI). France: UNESCO.
- [58] Wang, H. J., Yang, Z. S., Saito, Y., Liu, J. P., Sun, X. X., & Wang, Y. (2007). Stepwise decreases of the Huanghe (Yellow River) sediment load (1950–2005): Impacts of climate change and human activities. *Global and Planetary Change*, 57, 331–354. <https://doi.org/10.1016/j.gloplacha.2007.01.003>
- [59] Xu, X. T., Wang, Y. B., Yin, Z. H., & Zhang, H. (2017). Effect of temperature and strain rate on mechanical characteristics and constitutive model of frozen Helin loess. *Cold Regions Science and Technology*, 136, 44–51. <https://doi.org/10.1016/j.coldregions.2017.01.010>
- [60] Zádorová, T., Penížek, V., Šefrna, L., Drábek, O., Mihaljevič, M., Volf, S., & Chuman, T. (2013). Identification of Neolithic to Modern erosion-sedimentation phases using geochemical approach in a loess covered sub-catchment of South Moravia, Czech Republic. *Geoderma*, 195–196, 56–69. <https://doi.org/10.1016/j.geoderma.2012.11.012>
- [61] Zgłobicki, W. (2013). Present and past sedimentation rates in loess areas of the Lublin Upland (E Poland). *Geomorphologie*, 19(1), 79–92. <https://doi.org/10.4000/geomorphologie.10140>
- [62] Zhao, G., Mu, X., Wen, Z., Wang, F., & Gao, P. (2013). Soil erosion, conservation, and eco-environment changes in the Loess Plateau of China. *Land Degradation & Development*, 24, 499–510. <https://doi.org/10.1002/ldr.2246>
- [63] Zolina, O. (2012). Change in intense precipitation in Europe. In Z. W. Kundzewicz (Ed.), *Changes in Flood Risk in Europe* (pp. 97–120). Wallingford: IAHS Press and CRC Press/Balkema (Taylor & Francis Group) IAHS Special Publ. No. 10.